

Treatment/Outcome Literature 2000-2023 (89)

Arnow, B. A., Manber, R., Blasey, C., Klein, D. N., Blalock, J. A., Markowitz, J. C., Rothbaum, B. O., Rush, A. J., Thase, M. E., Riso, L. P., Vivian, D., McCullough, J. P. Jr., & Keller, M. B. (2003). Therapeutic Reactance as a Predictor of Outcome in the Treatment of Chronic Depression. *Journal of Consulting and Clinical Psychology*, 71(6), 1025–1035.
<https://doi.org/10.1037/0022-006X.71.6.1025>

Arnow, B. A., Blasey, C., Manber, R., Constantino, M. J., Markowitz, J. C., Klein, D. N., Thase, M. E., Kocsis, J. H., & Rush, A. J. (2007). Dropouts versus completers among chronically depressed outpatients. *Journal of Affective Disorders*, 97(1–3), 197–202.
<https://www.sciencedirect.com/science/article/pii/S0165032706002850>

Arnow, B. A., Steidtmann, D., Blasey, C., Manber, R., Constantino, M. J., Klein, D. N., Markowitz, J. C., Rothbaum, B. O., Thase, M. E., Fisher, A. J., & Kocsis, J. H. (2013). The relationship between the therapeutic alliance and treatment outcome in two distinct psychotherapies for chronic depression. *Journal of Consulting and Clinical Psychology*, 81(4), 627–638. <https://doi.org/10.1037/a0031530>

Assmann, N., Schramm, E., Kriston, L., Hautzinger, M., Härtter, M., Schweiger, U., & Klein, J. P. (2018). Moderating effect of comorbid anxiety disorders on treatment outcome in a randomized controlled psychotherapy trial in early-onset persistently depressed outpatients. *Depression & Anxiety*, 35(10), 1001–1008. <https://doi.org/10.1002/da.22839>

Bagean Koulemarz, M., Karami, J., Momeni, K., & Elahi, A. (2019). The effectiveness of cognitive behavioral analysis system of psychotherapy (CBASP) and integration of dialectical behavior therapy and cognitive therapy based on compassionate-mind on reduction mental pain and difficult emotion regulation people attempted suicide. *Clinical Psychology Studies*, 9(34), 35–63. https://jcps.atu.ac.ir/article_9879_en.html

Bagian Kulehmarzi, M. J., Karami, J., Momeni, K., & Elahi, A. (2018). The effectiveness of cognitive behavioral analysis system of psychotherapy (CBASP) and dialectical behavior therapy based on compassion on resilience and motivations for suicide attempts in people attempted suicide. *Contemporary Psychology, Biannual Journal of the Iranian Psychological Association*, 13(1), 26–39. http://bjcp.ir/browse.php?a_id=1376&sid=1&slc_lang=en

Bausch, P., Fangmeier, T., Meister, R., Elsaesser, M., Kriston, L., Klein, J. P., Zobel, I., Hautzinger, M., Härtter, M., & Schramm, E. (2020). The impact of childhood maltreatment on long-term outcomes in Disorder-Specific vs. Nonspecific psychotherapy for chronic depression. *Journal of Affective Disorders*, 272, 152–157.
<https://www.sciencedirect.com/science/article/pii/S0165032719321536>

Bausch, P., Fangmeier, T., Zobel, I., Schoepf, D., Drost, S., Schnell, K., Walter, H., Berger, M., Normann, C., & Schramm, E. (2017). The impact of childhood maltreatment on the differential efficacy of CBASP versus escitalopram in patients with chronic depression: A secondary analysis. *Clinical Psychology & Psychotherapy*, 24(5), 1155–1162.
<https://doi.org/10.1002/cpp.2081>

Berger, M., Brakemeier, E.-L., & Schramm, E. (2011). CS05-04-Cognitive behavioral analysis system of psychotherapy (CBASP). *European Psychiatry*, 26, 1785.
<https://www.sciencedirect.com/science/article/pii/S0924933811734897>

Blalock, J. A., Minnix, J. A., Mathew, A. R., Wetter, D. W., McCullough, J. P. Jr., & Cinciripini, P. M. (2013). Relationship of childhood trauma to depression and smoking outcomes in pregnant smokers. *Journal of Consulting and Clinical Psychology*, 81(5), 821–830. <https://doi.org/10.1037/a0033381>

Bollmann, S., Spies, J., Hodzic, J., Hohl-Radke, F., & Brakemeier, E.-L. (2020). Feasibility and outcome of an inpatient interpersonal skills group for older adults (65+) with acute or chronic depression: An open pilot study. *GeroPsych: The Journal of Gerontopsychology and Geriatric Psychiatry*, 33(2), 55–66. <https://doi.org/10.1024/1662-9647/a000224>

Brakemeier, E.-L., & Berger, M. (2011). ECP07-01-Cbasp workshop-introduction into the cognitive-behavioral analysis system of psychotherapy (cbasp) to treat depression. *European Psychiatry*, 26(S2), 1804–1804. <https://www.cambridge.org/core/journals/european-psychiatry/article/ecp0701-cbasp-workshop-introduction-into-the-cognitive-behavioral-analysis-system-of-psychotherapy-cbasp-to-treat-depression/583BC1151349BC32665368A81A7F7A71>

Brakemeier, E.-L., Engel, V., Schramm, E., Zobel, I., Schmidt, T., Hautzinger, M., Berger, M., & Normann, C. (2011). Feasibility and outcome of Cognitive Behavioral Analysis System of Psychotherapy (CBASP) for chronically depressed inpatients. *Psychotherapy and Psychosomatics*, 80(3), 191–194. <https://www.jstor.org/stable/48511340>

Brakemeier, E.-L., Herzog, P., Radtke, M., Schneibel, R., Breger, V., Becker, M., Spies, J., Jacobi, F., Heider, J., & Normann, C. (2018). CBASP as an Inpatient Concept for Treatment-Resistant Chronic Depression: A Pilot Study of the Relationship between Adverse Effects and Treatment Outcome. *Psychotherapie, Psychosomatik, Medizinische Psychologie*, 68(9–10), 399–407. <https://doi.org/10.1055/a-0629-7802>

Carney, C. E., Segal, Z. V., Edinger, J. D., & Krystal, A. D. (2007). A comparison of rates of residual insomnia symptoms following pharmacotherapy or cognitive-behavioral therapy for major depressive disorder. *Journal of Clinical Psychiatry*, 68(2), 254–260.
<https://www.psychiatrist.com/read-pdf/3503/>

Cinciripini, P. M., Blalock, J. A., Minnix, J. A., Robinson, J. D., Brown, V. L., Lam, C., Wetter, D. W., Schreindorfer, L., McCullough, J. P. Jr., Dolan-Mullen, P., Stotts, A. L., & Karam-Hage, M. (2010). Effects of an intensive depression-focused intervention for smoking cessation in pregnancy. *Journal of Consulting and Clinical Psychology*, 78(1), 44–54. <https://doi.org/10.1037/a0018168>

Constantino, M. J., Laws, H. B., Arnow, B. A., Klein, D. N., Rothbaum, B. O., & Manber, R. (2012). The relation between changes in patients' interpersonal impact messages and outcome in treatment for chronic depression. *Journal of Consulting and Clinical Psychology*, 80(3), 354–364. <https://doi.org/10.1037/a0028351>

Constantino, M. J., Laws, H. B., Coyne, A. E., Greenberg, R. P., Klein, D. N., Manber, R., Rothbaum, B. O., & Arnow, B. A. (2016). Change in patients' interpersonal impacts as a mediator of the alliance-outcome association in treatment for chronic depression. *Journal of Consulting and Clinical Psychology*, 84(12), 1135–1144. <https://doi.org/10.1037/ccp0000149>

Constantino, M. J., Manber, R., DeGeorge, J., McBride, C., Ravitz, P., Zuroff, D. C., Klein, D. N., Markowitz, J. C., Rothbaum, B. O., Thase, M. E., & Arnow, B. A. (2008). Interpersonal styles of chronically depressed outpatients: Profiles and therapeutic change: Psychotherapy. *Psychotherapy: Theory, Research, Practice, Training*, 45(4), 491–506. <https://doi.org/10.1037/a0014335>

Cukrowicz, K. C. (2005). *Prevention of anxiety and depression*. <https://diginole.lib.fsu.edu/islandora/object/fsu:181164>

Dean, J. A., Eldering, M. J., Schoevers, R. A., & van Driel, C. M. G. (2024). Identifying predictors of a favourable outcome for outpatients with a persistent depressive disorder treated with cognitive behavioural analysis system of psychotherapy: A prospective cohort study. *British Journal of Clinical Psychology*. <https://doi.org/10.1111/bjc.12454>

Denton, W. H., Carmody, T. J., Rush, A. J., Thase, M. E., Trivedi, M. H., Arnow, B. A., Klein, D. N., & Keller, M. B. (2010). Dyadic discord at baseline is associated with lack of remission in the acute treatment of chronic depression. *Psychological Medicine*, 40(3), 415–424. <https://www.cambridge.org/core/journals/psychological-medicine/article/dyadic-discord-at-baseline-is-associated-with-lack-of-remission-in-the-acute-treatment-of-chronic-depression/A8DC14824224DAD1A4ECFC85A786D057>

Echterhoff, J., Kriston, L., Klein, J. P., Härter, M., Schramm, E., & Schumacher, L. (2024). Symptom-specific improvement across therapies and their putative mediators: A mediation network intervention analysis. *Psychotherapy Research*. <https://doi.org/10.1080/10503307.2024.2320349>

Eich, H. S. (2019). *The Impact of Childhood Maltreatment on the Alliance in the Treatment for Persistent Depression: Analyses on the Differential Effects of the Cognitive Behavioral Analysis System of Psychotherapy and Supportive Psychotherapy* [PhD Thesis]. <https://archiv.ub.uni-heidelberg.de/volltextserver/26124/>

Emmelkamp, J., Kooistra, L., Van Oppen, P., Van Schaik, D., Hoogendoorn, A., Dekker, J., Beekman, A., McCullough, J. Jr., & Wiersma, J. (2020). Long-term outcome of treatment for persistent depressive disorder: A naturalistic follow-up study. *Psychotherapy and Psychosomatics*, 89(1), 54–55. <https://doi.org/10.1159/000503115>

Erkens, N., Schramm, E., Kriston, L., Hautzinger, M., Härter, M., Schweiger, U., & Klein, J. P. (2018). Association of comorbid personality disorders with clinical characteristics and outcome in a randomized controlled trial comparing two psychotherapies for early-onset persistent depressive disorders. *Journal of Affective Disorders*, 229, 262–268. <https://doi.org/10.1016/j.jad.2017.12.091>

Forkmann, T., Brakemeier, E.-L., Teismann, T., Schramm, E., & Michalak, J. (2016). The effects of Mindfulness-based Cognitive Therapy and Cognitive Behavioral Analysis System of Psychotherapy added to treatment as usual on suicidal ideation in chronic depression: Results of a randomized-clinical trial. *Journal of Affective Disorders*, 200, 51–57. <https://doi.org/10.1016/j.jad.2016.01.047>

Furukawa, T. A., Schramm, E., Weitz, E. S., Salanti, G., Efthimiou, O., Michalak, J., Watanabe, N., Cipriani, A., Keller, M. B., Kocsis, J. H., Klein, D. N., & Cuijpers, P. (2016). Cognitive-Behavioural Analysis System of Psychotherapy (CBASP), a drug, or their combination: Differential therapeutics for persistent depressive disorder: a study protocol of an individual participant data network meta-analysis. *BMJ Open*, 6(5), e011769. <https://doi.org/10.1136/bmjopen-2016-011769>

Ghadampour, E., & Bavazin, F. (2022). The Effectiveness of Cognitive-Behavioral Analysis Psychotherapy System (CBASP) on Attentional Bias in Adolescent Students with Symptoms of Social Anxiety. *Journal of Psychological Studies*, 17(4), 57–77.

Glanert, S., Sürig, S., Grave, U., Fassbinder, E., Schwab, S., Borgwardt, S., & Klein, J. P. (2021). Investigating care dependency and its relation to outcome (ICARE): Results from a naturalistic study of an intensive day treatment program for depression. *Frontiers in Psychiatry*, 12. <https://doi.org/10.3389/fpsy.2021.644972>

Guhn, A., Köhler, S., Brakemeier, E.-L., & Sterzer, P. (2021). Cognitive Behavioral Analysis System of Psychotherapy for inpatients with persistent depressive disorder: A naturalistic trial on a general acute psychiatric unit. *European Archives of Psychiatry & Clinical Neuroscience*, 271(3), 495–505. <https://doi.org/10.1007/s00406-019-01038-5>

Guhn, A., Merkel, L., Heim, C., Klawitter, H., Teich, P., Betzler, F., Sterzer, P., & Köhler, S. (2022). Impaired empathic functioning in chronic depression: Behavioral evidence for the Cognitive Behavioral Analysis System of Psychotherapy (CBASP) model. *Journal of Psychiatric Research*, 152, 79–85. <https://www.sciencedirect.com/science/article/pii/S0022395622003260>

Guhn, A., Schön, D., Zische, Y., Sterzer, P., & Köhler, S. (2021). Interpersonal change during inpatient CBASP treatment: Focus on group therapy. *Frontiers in Psychiatry*, 12, 620037.

<https://www.frontiersin.org/journals/psychiatry/articles/10.3389/fpsy.2021.620037/full>

Hirschfeld, R. M., Dunner, D. L., Keitner, G., Klein, D. N., Koran, L. M., Kornstein, S. G., Markowitz, J. C., Miller, I., Nemeroff, C. B., & Ninan, P. T. (2002). Does psychosocial functioning improve independent of depressive symptoms? A comparison of nefazodone, psychotherapy, and their combination. *Biological Psychiatry*, 51(2), 123–133.

<https://www.sciencedirect.com/science/article/pii/S0006322301012914>

Hom, M. A., Stanley, I. H., Vazquez, A., Belz, M., & Joiner, T. E. (2017). Gains in cognitive behavioral analysis system of psychotherapy: Examining treatment progress and processes of change in a psychiatric outpatient sample. *International Journal of Cognitive Therapy*, 10(3), 255–268. <https://doi.org/10.1521/ijct.2017.10.3.255>

Humer, E., Schramm, E., Klein, J. P., Härter, M., Hautzinger, M., Pieh, C., & Probst, T. (2021). Effects of alliance ruptures and repairs on outcomes. *Psychotherapy Research*, 31(8), 977–987. <https://doi.org/10.1080/10503307.2021.1874070>

Keller, M. B., McCullough, J. P., Klein, D. N., Arnow, B., Dunner, D. L., Gelenberg, A. J., Markowitz, J. C., Nemeroff, C. B., Russell, J. M., Thase, M. E., Trivedi, M. H., & Zajecka, J. (2000). A comparison of nefazodone, the cognitive behavioral-analysis system of psychotherapy, and their combination for the treatment of chronic depression. *The New England Journal of Medicine*, 342(20), 1462–1470.

<https://doi.org/10.1056/NEJM200005183422001>

Klein, D. N., Leon, A. C., Li, C., D'Zurilla, T. J., Black, S. R., Vivian, D., Dowling, F., Arnow, B. A., Manber, R., & Markowitz, J. C. (2011). Social problem solving and depressive symptoms over time: A randomized clinical trial of cognitive-behavioral analysis system of psychotherapy, brief supportive psychotherapy, and pharmacotherapy. *Journal of Consulting and Clinical Psychology*, 79(3), 342. <https://psycnet.apa.org/record/2011-07796-001>

Klein, D. N., Santiago, N. J., Vivian, D., Blalock, J. A., Kocsis, J. H., Markowitz, J. C., McCullough, J. P. Jr., Rush, A. J., Trivedi, M. H., Arnow, B. A., Dunner, D. L., Manber, R., Rothbaum, B., Thase, M. E., Keitner, G. I., Miller, I. W., & Keller, M. B. (2004). Cognitive-Behavioral Analysis System of Psychotherapy as a Maintenance Treatment for Chronic Depression. *Journal of Consulting and Clinical Psychology*, 72(4), 681–688.

<https://doi.org/10.1037/0022-006X.72.4.681>

Klein, D. N., Schwartz, J. E., Santiago, N. J., Vivian, D., Vocisano, C., Castonguay, L. G., Arnow, B., Blalock, J. A., Manber, R., & Markowitz, J. C. (2003). Therapeutic alliance in depression treatment: Controlling for prior change and patient characteristics. *Journal of Consulting and Clinical Psychology*, 71(6), 997. <https://psycnet.apa.org/record/2003-09784-005>

Klein, J. P., Becker, B., Hurlemann, R., Scheibe, C., Colla, M., & Heuser, I. (2014a). Effect of specific psychotherapy for chronic depression on neural responses to emotional faces. *Journal of Affective Disorders*, 166, 93–97.

<https://www.sciencedirect.com/science/article/pii/S0165032714002493>

Klein, J. P., Erkens, N., Schweiger, U., Kriston, L., Bausch, P., Zobel, I., Hautzinger, M., Schoepf, D., Serbanescu, I., & Bailer, J. (2018). Does childhood maltreatment moderate the effect of the cognitive behavioral analysis system of psychotherapy versus supportive psychotherapy in persistent depressive disorder? *Psychotherapy and Psychosomatics*, 87(1), 46–48. <https://karger.com/pps/article/87/1/46/283042>

Klein, J. P., Probst, T., Kriston, L., Assmann, N., Bailer, J., Eich, H., Schweiger, U., Stenzel, N. M., Wambach, K., Hautzinger, M., Härter, M., & Schramm, E. (2020). Changes in therapeutic alliance and in social inhibition as mediators of the effect of the cognitive behavioral analysis system of psychotherapy: A secondary analysis from a randomized clinical trial. *Psychotherapy and Psychosomatics*, 89(4), 261–262.

<https://doi.org/10.1159/000506082>

Knies, E.-M., Backenstrass, M., & Brakemeier, E.-L. (2016). Feasibility and outcome of a short-term group therapy for inpatients with comorbid substance use disorders and depression: The InterPersonal Kiesler-Circle Training (IPKCT). *Psychotherapy and Psychosomatics*, 85(5), 311–313. <https://doi.org/10.1159/000446157>

Kocsis, J. H., Gelenberg, A. J., Rothbaum, B. O., Klein, D. N., Trivedi, M. H., Manber, R., Keller, M. B., Leon, A. C., Wisniewski, S. R., Arnow, B. A., Markowitz, J. C., & Thase, M. E. (2009). Cognitive behavioral analysis system of psychotherapy and brief supportive psychotherapy for augmentation of antidepressant nonresponse in chronic depression: The REVAMP trial. *Archives of General Psychiatry*, 66(11), 1178–1188.

<https://doi.org/10.1001/archgenpsychiatry.2009.144>

Kocsis, J. H., Leon, A. C., Markowitz, J. C., Manber, R., Arnow, B., Klein, D. N., & Thase, M. E. (2009). Patient preference as a moderator of outcome for chronic forms of major depressive disorder treated with nefazodone, cognitive behavioral analysis system of psychotherapy, or their combination. *The Journal of Clinical Psychiatry*, 70(3), 354–361.

<https://doi.org/10.4088/JCP.08m04371>

Kocsis, J. H., Rush, A. J., Markowitz, J. C., Borian, F. E., Dunner, D. L., Koran, L. M., Klein, D. N., Trivedi, M. H., Arnow, B., Keitner, G., Kornstein, S. G., & Keller, M. B. (2003). Continuation treatment of chronic depression: A comparison of nefazodone, cognitive behavioral analysis system of psychotherapy, and their combination. *Psychopharmacology Bulletin*, 37(4), 73–87.

Konvalin, F., Grosse-Wentrup, F., Nenov-Matt, T., Fischer, K., Barton, B. B., Goerigk, S., Brakemeier, E.-L., Musil, R., Jobst, A., Padberg, F., & Reinhard, M. A. (2021). Borderline personality features in patients with persistent depressive disorder and their effect on CBASP outcome. *Frontiers in Psychiatry*, 12. <https://doi.org/10.3389/fpsyg.2021.608271>

Linstead, J., & Doyle, M. (2022). Evaluation of the impact and acceptability of Cognitive Behavioural Analysis System of Psychotherapy (CBASP) for chronic depression. *Behavioural & Cognitive Psychotherapy*, 50(6), 644–648. <https://doi.org/10.1017/S1352465822000236>

Locke, K. D., Sayegh, L., Penberthy, J. K., Weber, C., Haentjens, K., & Turecki, G. (2017). Interpersonal Circumplex Profiles Of Persistent Depression: Goals, Self-Efficacy, Problems, And Effects Of Group Therapy. *Journal of Clinical Psychology*, 73(6), 595–611. <https://doi.org/10.1002/jclp.22343>

Manber, R., Rush, A. J., Thase, M. E., Arnow, B., Klein, D., Trivedi, M. H., Korenstein, S. G., Markowitz, J. C., Dunner, D. L., Munsaka, M., Borian, F. E., & Keller, M. B. (2003). The effects of psychotherapy, nefazodone, and their combination on subjective assessment of disturbed sleep in chronic depression. *Sleep: Journal of Sleep and Sleep Disorders Research*, 26(2), 130–136. <https://doi.org/10.1093/sleep/26.2.130>

Meister, R., Lanio, J., Fangmeier, T., Härter, M., Schramm, E., Zobel, I., Hautzinger, M., Nestoriuc, Y., & Kriston, L. (2020). Adverse events during a disorder-specific psychotherapy compared to a nonspecific psychotherapy in patients with chronic depression. *Journal of Clinical Psychology*, 76(1), 7–19. <https://doi.org/10.1002/jclp.22869>

Michalak, J., Probst, T., Heidenreich, T., Bissantz, N., & Schramm, E. (2016). Mindfulness-based cognitive therapy and a group version of the cognitive behavioral analysis system of psychotherapy for chronic depression: Follow-up data of a randomized controlled trial and the moderating role of childhood adversities. *Psychotherapy and Psychosomatics*, 85(6), 378–380. <https://doi.org/10.1159/000447014>

Michalak, J., Schultze, M., Heidenreich, T., & Schramm, E. (2015). A randomized controlled trial on the efficacy of mindfulness-based cognitive therapy and a group version of cognitive behavioral analysis system of psychotherapy for chronically depressed patients. *Journal of Consulting and Clinical Psychology*, 83(5), 951. <https://psycnet.apa.org/record/2015-36864-001>

Negt, P., Brakemeier, E.-L., Michalak, J., Winter, L., Bleich, S., & Kahl, K. G. (2016). The treatment of chronic depression with cognitive behavioral analysis system of psychotherapy: A systematic review and meta-analysis of randomized-controlled clinical trials. *Brain and Behavior*, 6(8), e00486. <https://doi.org/10.1002/brb3.486>

Ninan, P. T., Rush, A. J., Crits-Christoph, P., Kornstein, S. G., Manber, R., Thase, M. E., Trivedi, M. H., Rothbaum, B. O., Zajecka, J., Borian, F. E., & Keller, M. B. (2002). Symptomatic and syndromal anxiety in chronic forms of major depression: Effect of nefazodone, Cognitive Behavioral Analysis System of Psychotherapy, and their combination. *The Journal of Clinical Psychiatry*, 63(5), 434–441. <https://doi.org/10.4088/JCP.v63n0510>

Parikh, S. V., Segal, Z. V., Grigoriadis, S., Ravindran, A. V., Kennedy, S. H., Lam, R. W., & Patten, S. B. (2009). Canadian Network for Mood and Anxiety Treatments (CANMAT) clinical guidelines for the management of major depressive disorder in adults II Psychotherapy alone or in combination with antidepressant medication. *Journal of Affective Disorders*, 117(Suppl 1), S15–S25. <https://doi.org/10.1016/j.jad.2009.06.042>

Penberthy, J. K., Gioia, C. J., Konig, A., Martin, A. M., Cockrell, S. A., & Meshberg-Cohen, S. (2014). Co-occurring chronic depression and alcohol dependence: A novel treatment approach. *Addictive Disorders & Their Treatment*, 13(2), 54–67. <https://doi.org/10.1097/ADT.0b013e31827de84e>

Probst, T., Schramm, E., Heidenreich, T., Klein, J., & Michalak, J. (2020). Patients' interpersonal problems as moderators of depression outcomes in a randomized controlled trial comparing mindfulness-based cognitive therapy and a group version of the cognitive-behavioral analysis system of psychotherapy in chronic depression. *Journal of Clinical Psychology*, 76(7), 1241–1254. <https://doi.org/10.1002/jclp.22931>

R. Manber, B. Arnow, C. Blasey, D. Vivian, J. P. McCULLOUGH, J. a. Blalock, D. N. Klein, J. C. Markowitz, L. P. Riso, B. Rothbaum, A. J. Rush, M. E. Thase, & M. B. Keller. (2003). Patient's therapeutic skill acquisition and response to psychotherapy, alone or in combination with medication. *Psychological Medicine*, 33(4), 693–702. <https://doi.org/10.1017/S0033291703007608>

Reuveni, I., Lauria, M., Monk, C., & Werner, E. (2021). The impact of childhood trauma on psychological interventions for depression during pregnancy and postpartum: A systematic review. *Archives of Women's Mental Health*, 24(3), 367–380. <https://doi.org/10.1007/s00737-020-01066-4>

Rief, W., Bleichhardt, G., Dannehl, K., Euteneuer, F., & Wambach, K. (2018). Comparing the efficacy of CBASP with two versions of CBT for depression in a routine care center: A randomized clinical trial. *Psychotherapy and Psychosomatics*, 87(3), 164–178. <https://doi.org/10.1159/000487893>

Rush, A. J., Trivedi, M. H., Carmody, T. J., Ibrahim, H. M., Markowitz, J. C., Keitner, G. I., Kornstein, S. G., Arnow, B., Klein, D. N., & Manber, R. (2005). Self-reported depressive symptom measures: Sensitivity to detecting change in a randomized, controlled trial of chronically depressed, nonpsychotic outpatients. *Neuropsychopharmacology*, 30(2), 405–416. <https://www.nature.com/articles/1300614>

Santiago, N. J., Klein, D. N., Vivian, D., Arnow, B. A., Blalock, J. A., Kocsis, J. H., Markowitz, J. C., Manber, R., Riso, L. P., Rothbaum, B. O., Rush, A. J., Thase, M. E., McCullough Jr., J. P., & Keller, M. B. (2005). The Therapeutic Alliance and CBASP-Specific Skill Acquisition in the Treatment of Chronic Depression. *Cognitive Therapy & Research*, 29(6), 803–817. <https://doi.org/10.1007/s10608-005-9638-5>

Santiago, N. J., Klein, D. N., Vivian, D., Vocisano, C., Dowling, F., Arnow, B. A., Manber, R., Markowitz, J. C., McCullough, J. P. Jr., Riso, L. P. L., Rothbaum, B. O., Rush, A. J., Thase, M. E., & Keller, M. B. (2002). Pretreatment correlates of the therapeutic alliance in the chronically depressed. *Journal of Contemporary Psychotherapy: On the Cutting Edge of Modern Developments in Psychotherapy*, 32(4), 281–290.
<https://doi.org/10.1023/A:1020524910971>

Satvat Qasriki, H., Hashemi Nosratabad, T., Bakhshi Pour Roudsari, A., & Mahmoud Alilou, M. (2023). Effectiveness of Cognitive Behavioral Analysis Psychotherapy System on Cognitive Reactivity and Reducing the Symptoms of Depressed Patients. *Journal of Research in Behavioural Sciences*, 20(4), 718–729.

http://rbs.mui.ac.ir/browse.php?a_id=1454&sid=1&slc_lang=en

Sayegh, L., Touré, E. H., Farquhar, E., Beaulieu, S., Renaud, S., Rej, S., & Perreault, M. (2020). Group Cognitive Behavioral Analysis System of Psychotherapy (CBASP): A pilot study for bipolar depression. *Frontiers in Psychiatry*, 11.

<https://doi.org/10.3389/fpsyg.2020.565681>

Schatzberg, A. F., Rush, A. J., Arnow, B. A., Banks, P. L. C., Blalock, J. A., Borian, F. E., Howland, R., Klein, D. N., Kocsis, J. H., Kornstein, S. G., Manber, R., Markowitz, J. C., Miller, I., Ninan, P. T., Rothbaum, B. O., Thase, M. E., Trivedi, M. H., & Keller, M. B. (2005). Chronic Depression: Medication (Nefazodone) or Psychotherapy (CBASP) Is Effective When the Other Is Not. *Archives of General Psychiatry*, 62(5), 513–520.

<https://doi.org/10.1001/archpsyc.62.5.513>

Schramm, E., Kriston, L., Elsaesser, M., Fangmeier, T., Meister, R., Bausch, P., Zobel, I., Bailer, J., Wambach, K., Backenstrass, M., Klein, J. P., Schoepf, D., Schnell, K., Gumz, A., Löwe, B., Walter, H., Wolf, M., Domschke, K., Berger, M., ... Härter, M. (2019). Two-year follow-up after treatment with the cognitive behavioral analysis system of psychotherapy versus supportive psychotherapy for early-onset chronic depression. *Psychotherapy and Psychosomatics*, 88(3), 154–164. <https://doi.org/10.1159/000500189>

Schramm, E., Kriston, L., Zobel, I., Bailer, J., Wambach, K., Backenstrass, M., Klein, J. P., Schoepf, D., Schnell, K., Gumz, A., Bausch, P., Fangmeier, T., Meister, R., Berger, M., Hautzinger, M., & Härter, M. (2017). Effect of disorder-specific vs nonspecific psychotherapy for chronic depression: A randomized clinical trial. *JAMA Psychiatry*, 74(3), 233–242. <https://doi.org/10.1001/jamapsychiatry.2016.3880>

Schramm, E., Zobel, I., Dykierk, P., Kech, S., Brakemeier, E.-L., Külz, A., & Berger, M. (2011). Cognitive behavioral analysis system of psychotherapy versus interpersonal psychotherapy for early-onset chronic depression: A randomized pilot study. *Journal of Affective Disorders*, 129(1–3), 109–116. <https://doi.org/10.1016/j.jad.2010.08.003>

Schramm, E., Zobel, I., Schoepf, D., Fangmeier, T., Schnell, K., Walter, H., Drost, S., Schmidt, P., Brakemeier, E.-L., Berger, M., & Normann, C. (2015). Cognitive behavioral analysis system of psychotherapy versus escitalopram in chronic major depression. *Psychotherapy and Psychosomatics*, 84(4), 227–240. <https://doi.org/10.1159/000381957>

Schumacher, L., Meister, R., & Kriston, L. (2022). *Using longitudinal symptom networks to assess symptom-specific effects of CBASP versus Supportive Psychotherapy for Chronic Depression*. <https://osf.io/dngra/resources>

Serbanescu, I., Backenstrass, M., Drost, S., Weber, B., Walter, H., Klein, J. P., Zobel, I., Hautzinger, M., Meister, R., Härter, M., Schramm, E., & Schoepf, D. (2020). Impact of baseline characteristics on the effectiveness of disorder-specific Cognitive Behavioral Analysis System of Psychotherapy (CBASP) and supportive psychotherapy in outpatient treatment for persistent depressive disorder. *Frontiers in Psychiatry*, 11. <https://doi.org/10.3389/fpsy.2020.607300>

Serbanescu, I., Schramm, E., Walter, H., Schnell, K., Zobel, I., Drost, S., Fangmeier, T., Normann, C., & Schoepf, D. (2024). Identifying subgroups with differential response to CBASP versus Escitalopram during the first eight weeks of treatment in outpatients with persistent depressive disorder. *European Archives of Psychiatry and Clinical Neuroscience*, 274(3), 723–737. <https://doi.org/10.1007/s00406-023-01672-0>

Serbanescu, I., Walter, H., Schnell, K., Kessler, H., Weber, B., Drost, S., Groß, M., Neudeck, P., Klein, J. P., Assmann, N., Zobel, I., Backenstrass, M., Hautzinger, M., Meister, R., Härter, M., Schramm, E., & Schoepf, D. (2020). Combining baseline characteristics to disentangle response differences to disorder-specific versus supportive psychotherapy in patients with persistent depressive disorder. *Behaviour Research and Therapy*, 124. <https://doi.org/10.1016/j.brat.2019.103512>

Shankman, S. A., Campbell, M. L., Klein, D. N., Leon, A. C., Arnow, B. A., Manber, R., Keller, M. B., Markowitz, J. C., Rothbaum, B. O., & Thase, M. E. (2013). Dysfunctional attitudes as a moderator of pharmacotherapy and psychotherapy for chronic depression. *Journal of Psychiatric Research*, 47(1), 113–121. <https://www.sciencedirect.com/science/article/pii/S0022395612002981>

Steidtmann, D., Manber, R., Blasey, C., Markowitz, J. C., Klein, D. N., Rothbaum, B. O., Thase, M. E., Kocsis, J. H., & Arnow, B. A. (2013). Detecting critical decision points in psychotherapy and psychotherapy+ medication for chronic depression. *Journal of Consulting and Clinical Psychology*, 81(5), 783. <https://psycnet.apa.org/record/2013-20187-001>

Swan, J. S., MacVicar, R., Christmas, D., Durham, R., Rauchhaus, P., McCullough, J. P. Jr., & Matthews, K. (2014). Cognitive Behavioural Analysis System of Psychotherapy (CBASP) for chronic depression: Clinical characteristics and six month clinical outcomes in an open case series. *Journal of Affective Disorders*, 152–154, 268–276. <https://doi.org/10.1016/j.jad.2013.09.024>

Szpak, K. (2020). *Does Cognitive Behavioural Analysis system of psychotherapy improve interpersonal functioning in patients affected by persistent depression and what can research tell us about the theoretical model this therapy is based on?*
<https://era.ed.ac.uk/handle/1842/37323>

Toussaint, A., Hüsing, P., Gumz, A., Wingenfeld, K., Härter, M., Schramm, E., & Löwe, B. (2020). Sensitivity to change and minimal clinically important difference of the 7-item Generalized Anxiety Disorder Questionnaire (GAD-7). *Journal of Affective Disorders*, 265, 395–401. <https://www.sciencedirect.com/science/article/pii/S0165032719313643>

Trivedi, M. H., Kocsis, J. H., Thase, M. E., Morris, D. W., Wisniewski, S. R., Leon, A. C., Gelenberg, A. J., Klein, D. N., Niederehe, G., Schatzberg, A. F., Ninan, P. T., & Keller, M. B. (2008). REVAMP - Research Evaluating the Value of Augmenting Medication with Psychotherapy: Rationale and design. *Psychopharmacology Bulletin*, 41(4), 5–33.

Vocisano, C., Klein, D. N., Arnow, B., Rivera, C., Blalock, J. A., Rothbaum, B., Vivian, D., Markowitz, J. C., Kocsis, J. H., Manber, R., Castonguay, L., Rush, A. J., Borian, F. E., McCullough, J. P., Kornstein, S. G., Riso, L. P., & Thase, M. E. (2004). Therapist Variables That Predict Symptom Change in Psychotherapy With Chronically Depressed Outpatients. *Psychotherapy: Theory, Research, Practice, Training*, 41(3), 255–265.
<https://doi.org/10.1037/0033-3204.41.3.255>

Wiersma, J. E., van Schaik, D. J. F., van Oppen, P., McCullough, J. P. Jr., Schoevers, R. A., Dekker, J. J., Blom, M. B. J., Maas, K., Smit, J. H., Penninx, B. W. J. H., & Beekman, A. T. F. (2008). Treatment of chronically depressed patients: A multisite randomized controlled trial testing the effectiveness of “Cognitive Behavioral Analysis System of Psychotherapy” (CBASP) for chronic depressions versus usual secondary care. *BMC Psychiatry*, 8. <https://doi.org/10.1186/1471-244X-8-18>

Wiersma, J. E., Van Schaik, D. J., Hoogendorn, A. W., Dekker, J. J., Van, H. L., Schoevers, R. A., Blom, M. B., Maas, K., Smit, J. H., & McCullough Jr, J. P. (2014). The effectiveness of the cognitive behavioral analysis system of psychotherapy for chronic depression: A randomized controlled trial. *Psychotherapy and Psychosomatics*, 83(5), 263–269.
<https://karger.com/pps/article-abstract/83/5/263/282868>

Wiersma, J. E., van Schaik, D. J., Hoogendorn, A. W., Schoevers, R. A., Dekker, J. J., Van, H. L., Blom, M. B., Maas, K., Smit, J. H., & McCullough Jr, J. P. (n.d.). *The Effectiveness of Cognitive Behavioral Analysis System of Psychotherapy for Chronic Depression: A Multisite Randomized Controlled Trial*. Retrieved April 24, 2024, from
<https://research.vumc.nl/ws/files/823301/chapter%206.pdf>

Wiersma, J. E., van Schaik, D. J., van Oppen, P., McCullough Jr, J. P., Schoevers, R. A., Dekker, J. J., Blom, M. B., Maas, K., Smit, J. H., & Penninx, B. W. (2008). A Multisite Randomized Controlled Trial testing the Effectiveness of Cognitive Behavioral Analysis System of Psychotherapy for Chronic Depression: Study protocol. *BMC Psychiatry*, 8, 18. <https://research.vumc.nl/ws/files/823299/chapter%205.pdf>

Wolf, J., Padberg, F., Nenov-Matt, T., Amann, B. L., Barton, B. B., Tang, J., Glessner, G., Brakemeier, E.-L., Jobst, A., Musil, R., & Reinhard, M. A. (2022). Suicidal behaviors are associated with loneliness and decrease during inpatient CBASP treatment for persistent depressive disorder. *Journal of Psychiatric Research*, 154, 139–144.
<https://doi.org/10.1016/j.jpsychires.2022.07.059>

Zajecka, J., Dunner, D. L., Gelenberg, A. J., Hirschfeld, R. M. A., Kornstein, S. G., Ninan, P. T., Rush, A. J., Thase, M. E., Trivedi, M., Arnow, B. A., Borian, F. E., Manber, R., & Keller, M. B. (2002). Sexual function and satisfaction in the treatment of chronic major depression with nefazodone, psychotherapy and their combination. *The Journal of Clinical Psychiatry*, 63(8), 709–716. <https://doi.org/10.4088/JCP.v63n0809>

Descriptive Literature 2000-2023 (74)

Arnow, B. A., & Constantino, M. J. (2003). Effectiveness of psychotherapy and combination treatment for chronic depression. *Journal of Clinical Psychology*, 59(8), 893–905.
<https://doi.org/10.1002/jclp.10181>

Arnow, B. A., Blasey, C., Manber, R., Constantino, M. J., Markowitz, J. C., Klein, D. N., Thase, M. E., Kocsis, J. H., & Rush, A. J. (2007). Dropouts versus completers among chronically depressed outpatients. *Journal of Affective Disorders*, 97(1–3), 197–202.
<https://doi.org/10.1016/j.jad.2006.06.017>

Arnow, B. A., Steidtmann, D., Blasey, C., Manber, R., Constantino, M. J., Klein, D. N., Markowitz, J. C., Rothbaum, B. O., Thase, M. E., Fisher, A. J., & Kocsis, J. H. (2013). The relationship between the therapeutic alliance and treatment outcome in two distinct psychotherapies for chronic depression. *Journal of Consulting and Clinical Psychology*, 81(4), 627–638. <https://doi.org/10.1037/a0031530>

Baty, P. (2000). Does nefazadone alone, the cognitive behavioral-analysis system of psychotherapy, or the combination of both work best for patients with chronic depression? *Journal of Family Practice*, 49(8), 687–687.
<http://proxy.library.stonybrook.edu/login?url=https://search.ebscohost.com/login.aspx?direct=true&db=pbh&AN=3459043&site=ehost-live&scope=site>

Bausch, P., Fangmeier, T., Schramm, E., Zobel, I., Drost, S., Schnell, K., Walter, H., Berger, M., Schoepf, D., & Normann, C. (2017). Cognitive behavioral analysis system of psychotherapy versus escitalopram in patients with chronic depression: Results from a naturalistic long-term follow-up. *Psychotherapy and Psychosomatics*, 86(5), 308–310.
<https://doi.org/10.1159/000477133>

Bird, T. (2016). *Do chronically depressed individuals exhibit a hostile-submissive interpersonal style and what is the process of change in Cognitive Behavioural Analysis System of Psychotherapy?* <https://era.ed.ac.uk/handle/1842/22083>

Blalock, J. A., Fouladi, R. T., Cinciripini, P. M., Markowitz, J. C., Klein, D. N., Rothbaum, B. O., Arnow, B. A., Manber, R., Riso, L. P., Sui, D., & McCullough Jr., J. P. (2008). Cognitive and Behavioral Mediators of Combined Pharmacotherapy and Psychotherapy of Chronic Depression. *Cognitive Therapy & Research*, 32(2), 197–211.
<https://doi.org/10.1007/s10608-006-9092-z>

Brakemeier, E.-L. (2011). Feasibility and effectiveness of cognitive behavioral analysis system of psychotherapy for chronically depressed inpatients: A pilot study. *European Psychiatry*, 26(S2), 2165–2165. <https://www.cambridge.org/core/journals/european-psychiatry/article/feasibility-and-effectiveness-of-cognitive-behavioral-analysis-system-of-psychotherapy-for-chronically-depressed-inpatients-a-pilot-study/366ABF15B0CDAB441461F12B5C2F8E92>

Brakemeier, E.-L., Engel, V., Schramm, E., Zobel, I., Schmidt, T., Hautzinger, M., Berger, M., & Normann, C. (2011). Feasibility and outcome of cognitive behavioral analysis system of psychotherapy (CBASP) for chronically depressed inpatients: A pilot study. *Psychotherapy and Psychosomatics*, 80(3), 191–194. <https://doi.org/10.1159/000320779>

Brakemeier, E.-L., Guhn, A., Stapel, S., Reinhard, M. A., & Padberg, F. (2023). Inpatient psychotherapy of depressive disorders: Options and challenges. *Der Nervenarzt*, 94(3), 213–224. <https://doi.org/10.1007/s00115-023-01448-w>

Brakemeier, E.-L., Radtke, M., Engel, V., Zimmermann, J., Tuschen-Caffier, B., Hautzinger, M., Schramm, E., Berger, M., & Normann, C. (2015). Overcoming treatment resistance in chronic depression: A pilot study on outcome and feasibility of the Cognitive Behavioral Analysis System of Psychotherapy as an inpatient treatment program. *Psychotherapy and Psychosomatics*, 84(1), 51–56. <https://doi.org/10.1159/000369586>

Constantino, M. J., Laws, H. B., Arnow, B. A., Klein, D. N., Rothbaum, B. O., & Manber, R. (2012). The relation between changes in patients' interpersonal impact messages and outcome in treatment for chronic depression. *Journal of Consulting and Clinical Psychology*, 80(3), 354–364. <https://doi.org/10.1037/a0028351>

Constantino, M. J., Laws, H. B., Coyne, A. E., Greenberg, R. P., Klein, D. N., Manber, R., Rothbaum, B. O., & Arnow, B. A. (2016). Change in patients' interpersonal impacts as a mediator of the alliance-outcome association in treatment for chronic depression. *Journal of Consulting and Clinical Psychology*, 84(12), 1135–1144. <https://doi.org/10.1037/ccp0000149>

Constantino, M. J., Manber, R., DeGeorge, J., McBride, C., Ravitz, P., Zuroff, D. C., Klein, D. N., Markowitz, J. C., Rothbaum, B. O., & Thase, M. E. (2008). Interpersonal styles of chronically depressed outpatients: Profiles and therapeutic change. *Psychotherapy: Theory, Research, Practice, Training*, 45(4), 491. <https://psycnet.apa.org/record/2008-19012-006>

Cuijpers, P., Ciharova, M., Quero, S., Miguel, C., Driessen, E., Harrer, M., Purgato, M., Ebert, D., & Karyotaki, E. (2022). The contribution of “individual participant data” meta-analyses of psychotherapies for depression to the development of personalized treatments: A systematic review. *Journal of Personalized Medicine*, 12(1), 93.
<https://www.mdpi.com/2075-4426/12/1/93>

Dean, J. A., Eldering, M. J., Schoevers, R. A., & van Driel, C. M. G. (2024). Identifying predictors of a favorable outcome for outpatients with a persistent depressive disorder treated with cognitive behavioral analysis system of psychotherapy: A prospective cohort study. *British Journal of Clinical Psychology*. <https://doi.org/10.1111/bjc.12454>

Dhingra, S., & Parle, M. (2011). Non-drug strategies in the management of depression: A comprehensive study of systematic review and meta-analysis of randomized controlled trials. *J Neurosci Behav Health*, 3(5), 66–73.

https://www.researchgate.net/profile/Sameer_Dhingra2/publication/228478292_Non-drug_strategies_in_the_management_of_depression_A_comprehensive_study_of_systematic_review_and_meta-analysis_of_randomised_controlled_trials/links/0deec53b6cbe9deb62000000/Non-drug-strategies-in-the-management-of-depression-A-comprehensive-study-of-systematic-review-and-meta-analysis-of-randomised-controlled-trials.pdf

Dippel, N., In-Albon, T., Schneider, S., Christiansen, H., & Brakemeier, E.-L. (2022). Investigating the feasibility and effectiveness of a modular treatment program for children and adolescents with depression and interpersonal problems: Study protocol of a quasi-experimental pilot feasibility trial (CBASP@YoungAge). *Pilot and Feasibility Studies*, 8(1), 145. <https://doi.org/10.1186/s40814-022-01091-3>

Dirmaier, J., Steinmann, M., Krattenmacher, T., Watzke, B., Barghaan, D., Koch, U., & Schulz, H. (2012). Non-pharmacological treatment of depressive disorders: A review of evidence-based treatment options. *Reviews on Recent Clinical Trials*, 7(2), 141–149.
<https://www.ingentaconnect.com/content/ben/rrct/2012/00000007/00000002/art00009>

DiSalvo, C. A., & McCullough, J. P. Jr. (2002). Treating a chronically depressed adolescent female using the cognitive behavioral analysis system of psychotherapy: A case study: Journal of Contemporary Psychotherapy. *On the Cutting Edge of Modern Developments in Psychotherapy*, 32(4), 273–280. <https://doi.org/10.1023/A:1020572826900>

Echterhoff, J., Kriston, L., Klein, J. P., Härter, M., Schramm, E., & Schumacher, L. (2023). *The effect of psychotherapy on the relations between treatment targets and persistent depressive disorder: A Bayesian network analysis*. <https://psyarxiv.com/n7mkv/>

Elsaesser, M., Feige, B., Kriston, L., Schumacher, L., Peifer, J., Hautzinger, M., Härter, M., & Schramm, E. (2023). *Longitudinal clusters and predictors of long-term trajectories in patients with early-onset chronic depression: Two years of follow-up after psychological treatment*. <https://osf.io/preprints/psyarxiv/h32kp/>

Emmelkamp, J., Kooistra, L., Van Oppen, P., Van Schaik, D., Hoogendoorn, A., Dekker, J., Beekman, A., McCullough Jr, J., & Wiersma, J. (2020). Long-term outcome of treatment for persistent depressive disorder: A naturalistic follow-up study. *Psychotherapy and Psychosomatics*, 89(1), 54–55. <https://karger.com/pps/article/89/1/54/289786>

Favorite, T. K., & Conrad, K. J. (2013). Group Intervention for Chronic Depression and PTSD. *Psychiatric Disorders—New Frontiers in Affective Disorders*, 269. https://books.google.com/books?hl=en&lr=lang_en|lang_it&id=NkueDwAAQBAJ&oi=fnd&pg=PA269&dq=CBASP+or+cognitive+behavioral+analysis+system+of+psychotherapy+&ots=0NBdaiLLOM&sig=97MJYtO2AD-N-Vpt_WU14yt8tAU

Flynn, H. A., & Himle, J. (2011). *Psychotherapy strategies for treatment resistant depression*. <https://psycnet.apa.org/record/2011-07396-010>

Groh, A., Jahn, K., Walter, M., Heck, J., Lichtinghagen, R., Janke, E., Westhoff, M. L. S., Deest, M., Frieling, H., Bleich, S., Kahl, K. G., & Heberlein, A. (2021). TNF- α Increase in a Cohort of Depressive Patients. *Disease Markers*, 2021, 8897421. <https://doi.org/10.1155/2021/8897421>

Guhn, A., Merkel, L., Heim, C., Klawitter, H., Teich, P., Betzler, F., Sterzer, P., & Köhler, S. (2022). Impaired empathic functioning in chronic depression: Behavioral evidence for the Cognitive Behavioral Analysis System of Psychotherapy (CBASP) model. *Journal of Psychiatric Research*, 152, 79–85. <https://doi.org/10.1016/j.jpsychires.2022.06.020>

Hellerstein, D. J. (2008). Review of Treating chronic depression with disciplined personal involvement: Cognitive behavioral analysis system of psychotherapy (CBASP): Psychotherapy. *Theory, Research, Practice, Training*, 45(3), 416–417. <https://doi.org/10.1037/a0013330>

Herzog, P., Häusler, S., Normann, C., & Brakemeier, E.-L. (2021). Negative Effects of a Multimodal Inpatient CBASP Program: Rate of Occurrence and Their Impact on Treatment Outcome in Chronic and Treatment-Resistant Depression. *Frontiers in Psychiatry*, 12, 575837. <https://doi.org/10.3389/fpsyg.2021.575837>

Hirschfeld, R. M. A., Dunner, D. L., Keitner, G., Klein, D. N., Koran, L. M., Kornstein, S. G., Markowitz, J. C., Miller, I., Nemeroff, C. B., Ninan, P. T., Rush, A. J., Schatzberg, A. F., Thase, M. E., Trivedi, M. H., Borian, F. E., Crits-Christoph, P., & Keller, M. B. (2002). Does psychosocial functioning improve independent of depressive symptoms? A comparison of nefazodone, psychotherapy, and their combination. *Biological Psychiatry*, 51(2), 123–133. [https://doi.org/10.1016/S0006-3223\(01\)01291-4](https://doi.org/10.1016/S0006-3223(01)01291-4)

Hom, M. A., Stanley, I. H., Vazquez, A., Belz, M., & Joiner, T. E. (2017). Gains in cognitive behavioral analysis system of psychotherapy: Examining treatment progress and processes of change in a psychiatric outpatient sample. *International Journal of Cognitive Therapy*, 10(3), 255–268. <https://doi.org/10.1521/ijct.2017.10.3.255>

Horn, G. L. (2012). An investigation into the effects of cognitive behavioral therapy on patients with chronic depression: A small case series. *Psychology Research and Behavior Management*, 5.

Jehle, P. J., & McCullough, J. P. Jr. (2002). Treatment of chronic major depression using the cognitive behavioral analysis system of psychotherapy (CBASP): Journal of Contemporary Psychotherapy. *Journal of Contemporary Psychotherapy: On the Cutting Edge of Modern Developments in Psychotherapy*, 32(4), 263–271. <https://doi.org/10.1023/A:1020520810062>

Jobst, A., Sabaß, L., Hall, D., Brücklmeier, B., Buchheim, A., Hall, J., Sarubin, N., Zill, P., Falkai, P., Brakemeier, E.-L., & Padberg, F. (2018). Oxytocin plasma levels predict the outcome of psychotherapy: A pilot study in chronic depression. *Journal of Affective Disorders*, 227, 206–213. <https://doi.org/10.1016/j.jad.2017.10.037>

Klein, J. P., Dale, R., Glanert, S., Grave, U., Surig, S., Zurowski, B., Borgwardt, S., Schweiger, U., Fassbinder, E., & Probst, T. (2022). Does childhood emotional abuse moderate the effect of Cognitive Behavioral Analysis System of Psychotherapy versus Meta-Cognitive Therapy in depression? A propensity score analysis on an observational study. *Journal of Affective Disorders*, 300, 71–75. <https://doi.org/10.1016/j.jad.2021.12.087>

Knies, E.-M., Backenstrass, M., & Brakemeier, E.-L. (2016). Feasibility and Outcome of a Short-Term Group Therapy for Inpatients with Comorbid Substance Use Disorders and Depression: The InterPersonal Kiesler-Circle Training (IPKCT). *Psychotherapy and Psychosomatics*, 85(5), 311–313. <https://karger.com/pps/article/85/5/311/282964>

Kocsis, J. H., Leon, A. C., Markowitz, J. C., Manber, R., Arnow, B., Klein, D. N., & Thase, M. E. (2009). Patient preference as a moderator of outcome for chronic forms of major depressive disorder treated with nefazodone, cognitive behavioral analysis system of psychotherapy, or their combination. *The Journal of Clinical Psychiatry*, 70(3), 354–361. <https://doi.org/10.4088/JCP.08m04371>

Köhler, S., Fischer, T., Brakemeier, E.-L., & Sterzer, P. (2015). Successful treatment of severe persistent depressive disorder with a sequential approach: Electroconvulsive therapy followed by cognitive behavioral analysis system of psychotherapy. *Psychotherapy and Psychosomatics*, 84(2), 127–128. <https://karger.com/pps/article/84/2/127/289421>

Köhler, S., Guhn, A., Betzler, F., Stiglmayr, C., Brakemeier, E.-L., & Sterzer, P. (2017). Therapeutic self-disclosure within DBT, schema therapy, and CBASP: Opportunities and challenges. *Frontiers in Psychology*, 8. <https://doi.org/10.3389/fpsyg.2017.02073>

Kriston, L., Wolff, A., Westphal, A., Hözel, L. P., & Härtter, M. (2014). Efficacy and Acceptability of Acute Treatments for Persistent Depressive Disorder: A Network Meta-Analysis. *Depression & Anxiety*, 31(8), 621–630. <https://doi.org/10.1002/da.22236>

La Rocco, D. (2017). Group Workbook for Treatment of Persistent Depression: Cognitive Behavioral Analysis System of Psychotherapy (CBASP) Therapist's Guide by Lilane Sayegh and J. Kim Penberthy, and: Group Workbook for Treatment of Persistent Depression: Cognitive Behavioral Analysis System of Psychotherapy (CBASP) Patient's Guide by Lilane Sayegh and J. Kim Penberthy. *Group*, 41(3), 267–270.

<https://muse.jhu.edu/pub/440/article/844401/summary>

Long, L. R., Foster, M., Burr, K., Penberthy, J. K., Baker, T. N., & McCullough, J. P. (2024). Pilot study dismantling the cognitive behavioral analysis system of psychotherapy: Identifying the active ingredients. *Cognitive Therapy and Research*.

<https://doi.org/10.1007/s10608-024-10467-z>

Maddux, R. E., Riso, L. P., Klein, D. N., Markowitz, J. C., Rothbaum, B. O., Arnow, B. A., Manber, R., Blalock, J. A., Keitner, G. I., & Thase, M. E. (2009). Select comorbid personality disorders and the treatment of chronic depression with nefazodone, targeted psychotherapy, or their combination. *Journal of Affective Disorders*, 117(3), 174–179.

<https://www.sciencedirect.com/science/article/pii/S0165032709000081>

Manber, R., Arnow, B., Blasey, C., Vivian, D., McCullough, J. P., Blalock, J. A., Klein, D. N., Markowitz, J. C., Riso, L. P., & Rothbaum, B. (2003). Patient's therapeutic skill acquisition and response to psychotherapy, alone or in combination with medication. *Psychological Medicine*, 33(4), 693–702.

<https://www.cambridge.org/core/journals/psychological-medicine/article/patients-therapeutic-skill-acquisition-and-response-to-psychotherapy-alone-or-in-combination-with-medication/C707779280CB56E147B065F0EE85C3CA>

Manber, R., Kraemer, H. C., Arnow, B. A., Trivedi, M. H., Rush, A. J., Thase, M. E., Rothbaum, B. O., Klein, D. N., Kocsis, J. H., & Gelenberg, A. J. (2008). Faster remission of chronic depression with combined psychotherapy and medication than with each therapy alone. *Journal of Consulting and Clinical Psychology*, 76(3), 459.

<https://psycnet.apa.org/record/2008-06469-010>

McCullough Jr, J. P. (2000). The case of Katrina: Skating on thin ice. *Cognitive and Behavioral Practice*, 7(4), 510–514.

<https://www.sciencedirect.com/science/article/pii/S107772290080067X>

McCullough, J. P. Jr., Lord, B. D., Conley, K. A., & Martin, A. M. (2010). A method for conducting intensive psychological studies with early-onset chronically depressed patients. *American Journal of Psychotherapy*, 64(4), 317–337.

<http://proxy.library.stonybrook.edu/login?url=https://search.ebscohost.com/login.aspx?direct=true&db=psyh&AN=2010-26373-001&site=ehost-live&scope=site>

McCullough, J. P., Lord, B. D., Martin, A. M., Conley, K. A., Schramm, E., & Klein, D. N. (2011). The Significant Other History: An Interpersonal-Emotional History Procedure Used with the Early-onset Chronically Depressed Patient. *American Journal of Psychotherapy (Association for the Advancement of Psychotherapy)*, 65(3), 225–248.

<https://doi.org/10.1176/appi.psychotherapy.2011.65.3.225>

McPherson, S., & Senra, H. (2022). Psychological treatments for persistent depression: A systematic review and meta-analysis of quality of life and functioning outcomes.

Psychotherapy, 59(3), 447. <https://psycnet.apa.org/record/2022-76312-001>

McPherson, S., Cairns, P., Carlyle, J., Shapiro, D. A., Richardson, P., & Taylor, D. (2005). The effectiveness of psychological treatments for treatment-resistant depression: A systematic review. *Acta Psychiatrica Scandinavica*, 111(5), 331–340.

<https://doi.org/10.1111/j.1600-0447.2004.00498.x>

Nemeroff, C. B., Heim, C. M., Thase, M. E., Klein, D. N., Rush, A. J., Schatzberg, A. F., Ninan, P. T., McCullough, J. P., Weiss, P. M., Dunner, D. L., Rothbaum, B. O., Kornstein, S., Keitner, G., & Keller, M. B. (2003). Differential responses to psychotherapy versus pharmacotherapy in patients with chronic forms of major depression and childhood trauma. *Proceedings of the National Academy of Sciences of the United States of America*, 100(24), 14293–14296. <https://doi.org/10.1073/pnas.2336126100>

Netter, A.-L., Beintner, I., & Brakemeier, E.-L. (2022). Adding an App-Based Intervention to the Cognitive Behavioral Analysis System of Psychotherapy in Routine Outpatient Psychotherapy Treatment: Proof-of-Concept Study. *JMIR Formative Research*, 6(8), e35482. <https://doi.org/10.2196/35482>

Niciu, M. J., Abdallah, C. G., Fenton, L. R., Fasula, M. K., Black, A., Anderson, G. M., & Sanacora, G. (2015). A history of early life parental loss or separation is associated with successful cognitive-behavioral therapy in major depressive disorder. *Journal of Affective Disorders*, 187, 241–244.

<https://www.sciencedirect.com/science/article/pii/S0165032715303906>

Öst, L.-G. (2008). Efficacy of the third wave of behavioral therapies: A systematic review and meta-analysis. *Behavior Research and Therapy*, 46(3), 296–321.

<https://doi.org/10.1016/j.brat.2007.12.005>

Penberthy, K. J., Khanna, S., Morgan Lynch, D., Chhabra, M. L. T., Xu, Y., & Gioia, C. (2016). *Effective Treatment for Co-Occurring Alcohol Use Disorder and Persistent Depression: A*.

https://www.researchgate.net/profile/Surbhi_Khanna/publication/312084932_Effective_Treatment_for_Co-Occurring_Alcohol_use_Disorder_and_Persistent_Depression_A_Case_Report/links/586e63f108ae6eb871bcfd77/Effective-Treatment-for-Co-Occurring-Alcohol-use-Disorder-and-Persistent-Depression-A-Case-Report.pdf

Potijk, M. R., aan het Rot, M., Parlevliet, F. M., Schoevers, R. A., & Eldering, M. J. (2020). Group Cognitive Behavioural Analysis System of Psychotherapy (CBASP) for persistently depressed outpatients: A retrospective chart review. *British Journal of Clinical Psychology*, 59(4), 552–564. <https://doi.org/10.1111/bjc.12266>

Riso, L. P., McCullough, J. P., & Blandino, J. (2003). Focus on Empirically Supported Methods: The Cognitive Behavioral Analysis System of Psychotherapy: A Promising Treatment for Clinical Depression. *The Scientific Review of Mental Health Practice: Objective Investigations of Controversial and Unorthodox Claims in Clinical Psychology, Psychiatry, and Social Work*. <https://psycnet.apa.org/record/2004-20405-007>

Sabaß, L., Padberg, F., Normann, C., Engel, V., Konrad, C., Helmle, K., Jobst, A., Worlitz, A., & Brakemeier, E.-L. (2018). Cognitive Behavioral Analysis System of Psychotherapy as group psychotherapy for chronically depressed inpatients: A naturalistic multicenter feasibility trial. *European Archives of Psychiatry & Clinical Neuroscience*, 268(8), 783–796. <https://doi.org/10.1007/s00406-017-0843-5>

Salmi, K. (2017). *Effect of CBASP therapy on the neurophysiological responses to pictures of neutral faces in depression: An event-related potential study* [Master's Thesis]. <https://jyx.jyu.fi/handle/123456789/55201>

Santiago, N. J., Klein, D. N., Vivian, D., Vocisano, C., Dowling, F., Arnow, B. A., Manber, R., Markowitz, J. C., McCullough, J. P. Jr., Riso, L. P. L., Rothbaum, B. O., Rush, A. J., Thase, M. E., & Keller, M. B. (2002). Pretreatment correlates of the therapeutic alliance in the chronically depressed. *Journal of Contemporary Psychotherapy: On the Cutting Edge of Modern Developments in Psychotherapy*, 32(4), 281–290. <https://doi.org/10.1023/A:1020524910971>

Sayegh, L., Locke, K. D., Pistilli, D., Penberthy, J. K., Chachamovich, E., McCullough, J. P. Jr., & Turecki, G. (2012). Cognitive behavioral analysis system of psychotherapy for treatment-resistant depression: Adaptation to a group modality. *Behavior Change*, 29(2), 97–108. <https://doi.org/10.1017/bec.2012.2>

Schramm, E., Hautzinger, M., Zobel, I., Kriston, L., Berger, M., & Härter, M. (2011). Comparative efficacy of the Cognitive Behavioral Analysis System of Psychotherapy versus Supportive Psychotherapy for early onset chronic depression: Design and rationale of a multisite randomized controlled trial. *BMC Psychiatry*, 11. <https://doi.org/10.1186/1471-244X-11-134>

Schramm, P. J., Zobel, I., Mönch, K., Schramm, E., & Michalak, J. (2016). Sleep quality changes in chronically depressed patients treated with mindfulness-based cognitive therapy or the cognitive behavioral analysis system of psychotherapy: A pilot study. *Sleep Medicine*, 17, 57–63. <https://www.sciencedirect.com/science/article/pii/S1389945715020237>

Segal, Z., Vincent, P., & Levitt, A. (2002). Efficacy of combined, sequential and crossover psychotherapy and pharmacotherapy in improving outcomes in depression. *Journal of Psychiatry and Neuroscience*, 27(4), 281–290. <https://www.jpn.ca/content/27/4/281.short>

Sperry, L. (2007). Dealing with the spiritual issues of chronically ill clients in the context of couples counseling: A unique application of cognitive behavioral analysis system of psychotherapy (CBASP). *The Family Journal*, 15(2), 183–187.
<https://doi.org/10.1177/1066480706298729>

Steidtmann, D., Manber, R., Arnow, B. A., Klein, D. N., Markowitz, J. C., Rothbaum, B. O., Thase, M. E., & Kocsis, J. H. (2012). Patient Treatment Preference as a Predictor of Response and Attrition in Treatment for Chronic Depression. *Depression & Anxiety*, 29(10), 896–905. <https://doi.org/10.1002/da.21977>

Stulz, N., Thase, M. E., Klein, D. N., Manber, R., & Crits-Christoph, P. (2010). Differential effects of treatments for chronic depression: A latent growth model reanalysis. *Journal of Consulting and Clinical Psychology*, 78(3), 409–419. <https://doi.org/10.1037/a0019267>

Sürig, S., Ohm, K., Grave, U., Glanert, S., Herzog, P., Fassbinder, E., Borgwardt, S., & Klein, J. P. (2021). Change in interpersonal and metacognitive skills during treatment with cognitive behavioral analysis system of psychotherapy and metacognitive therapy: Results from an observational study. *Frontiers in Psychiatry*, 12.
<https://doi.org/10.3389/fpsy.2021.619674>

Sürig, S., Ohm, K., Grave, U., Glanert, S., Herzog, P., Fassbinder, E., Borgwardt, S., & Klein, J. P. (2022). Interpersonal skills deficits and dysfunctional metacognitive beliefs have been implicated in the etiology and maintenance of depression. *CBASP in the Treatment of Persistent Depressive Disorder*.
https://books.google.com/books?hl=en&lr=lang_en|lang_it&id=y9pbEAAAQBAJ&oi=fnd&pg=PA139&dq=CBASP+or+Cognitive+Behavioral+Analysis+System+of+Psychotherapy+&ots=EF4_UWJO0c&sig=rUzrWxI6QedFj4_I8Lt_tBoZRIE

van Bronswijk, S., Moopen, N., Beijers, L., Ruhe, H. G., & Peeters, F. (2019). Effectiveness of psychotherapy for treatment-resistant depression: A meta-analysis and meta-regression. *Psychological Medicine*, 49(3), 366–379. <https://doi.org/10.1017/S003329171800199X>

van Hees, M. L. J. M., Rotter, T., Ellermann, T., & Evers, S. M. A. A. (2013). The effectiveness of individual interpersonal psychotherapy as a treatment for major depressive disorder in adult outpatients: A systematic review: BMC Psychiatry. *BMC Psychiatry*, 13.
<https://doi.org/10.1186/1471-244X-13-22>

Vivian, D., & Salwen, J. (2013). Key process issues in Cognitive Behavioral Analysis System of Psychotherapy (CBASP): Translation of an evidence-based model into clinical practice and training. *Psychotherapy*, 50(3), 398. <https://psycnet.apa.org/record/2013-30487-028>

Winter, L., Kahl, K. G., Wells, A., & Schweiger, U. (2012). Individual psychotherapy for patients with major depression: New methods for new therapies. *Nervenheilkunde*, 31(5), 316–320. <https://research.manchester.ac.uk/en/publications/individual-psychotherapy-for-patients-with-major-depression-new-m>

Wolf, J., Padberg, F., Nenov-Matt, T., Amann, B. L., Barton, B. B., Tang, J., Glessner, G., Brakemeier, E.-L., Jobst, A., Musil, R., & Reinhard, M. A. (2022). Suicidal behaviors are associated with loneliness and decrease during inpatient CBASP treatment for persistent depressive disorder. *Journal of Psychiatric Research*, 154, 139–144. <https://doi.org/10.1016/j.jpsychires.2022.07.059>

Books, Manuals and Editorial Literature 1995-2023 (86)

Alavi, K. (2020). The Theory of Psychopathology and the Treatment Structure in Cognitive Behavioral Analysis System of Psychotherapy (CBASP) and it's Effectiveness on Chronic Depression. *Rooyesh-e-Ravanshenasi Journal (RRJ)*, 9(5), 193–202. http://frooyesh.ir/browse.php?a_id=1892&slc_lang=en&sid=1&printcase=1&hbnr=1&hmb=1

Arnow, B. A. (2005). Cognitive Behavioral Analysis System of Psychotherapy for Chronic Depression. *Cognitive and Behavioral Practice*, 12(1), 6–16. [https://doi.org/10.1016/S1077-7229\(05\)80035-5](https://doi.org/10.1016/S1077-7229(05)80035-5)

Berthold-Losleben, M., Liebing-Wilson, M., & Swan, J. S. (2023). *The ABCs of CBASP: A guide to the cognitive behavioral analysis system of psychotherapy for therapists and supervisors* (2023-65873-000). Hogrefe. <https://doi.org/10.1027/00584-000>

Bland, R. C. (2001). Treatment for chronic depression: Cognitive behavioral analysis system of psychotherapy (CBASP). *Journal of Psychiatry and Neuroscience*, 26(4), 342. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC167192/>

Bollmann, S., Radtke, M., Normann, C., Zimmermann, J., Kessler, E. M., & Brakemeier, E. L. (2015). The Cognitive Behavioral Analysis System of Psychotherapy (CBASP) for chronically depressed elderly patients. *Psychotherapy and Psychosomatics*, 84, 8–8.

Burt, V. K. (2004). Plotting the course to remission: The search for better outcomes in the treatment of depression. *Journal of Clinical Psychiatry*, 65, 20–25. https://www.psychiatrist.com/wp-content/uploads/2021/02/16076_plotting-course-remission-search-better-outcomes-treatment.pdf

Cardemil, E. V. (2014). Gabor I. Keitner Brown University and Rhode Island Hospital Providence, Rhode Island, USA Esteban V. Cardemil Clark University. *Handbook of Chronic Depression: Diagnosis and Therapeutic Management*, 25, 134.
https://books.google.com/books?hl=en&lr=lang_en|lang_it&id=KFPVPlzuOX88C&oi=fnd&pg=PA134&dq=CBASP+or+cognitive+behavioral+analysis+system+of+psychotherapy+&ots=NnJlbzWPcW&sig=TplyS-jOSmOdQ9Gpu5CcjcDjv8I

Claiborn, J. M. (2001). Treatment for Chronic Depression: Cognitive Behavioral Analysis System of Psychotherapy. *The Journal of Psychotherapy Practice and Research*, 10(2), 136.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3330637/>

Coupland, J., Morrison, J., Myrh, G., Pinard, G., Racine, H., Tagalakis, V., St-Laurent, M., & Talbot, J. (2008). "New developments in cognitive-behavioural therapy for mood disorders". *The Canadian Journal of Psychiatry / La Revue Canadienne de Psychiatrie*, 53(6), 408–408.

Cox, K. C. (2011). *An Educational Workshop: Introducing an Evidence-Based Psychotherapy Strategy" Tool Kit" for the Treatment of Chronic Depression*.
<https://repository.usfca.edu/cgi/viewcontent.cgi?article=1009&context=dnp>

McCullough, J. P. (2006). Creating contingent environments using contingent personal responsiveness. *Treating Chronic Depression with Disciplined Personal Involvement* (pp. 81–122). Springer US. https://doi.org/10.1007/978-0-387-31066-4_5

Cuijpers, P., Huibers, M. J. H., & Furukawa, T. A. (2017). The need for research on treatments of chronic depression. *JAMA Psychiatry*, 74(3), 242–243.
<https://doi.org/10.1001/jamapsychiatry.2016.4070>

Curran, J., Houghton, S., & Grant, A. (2010). Developments within the Cognitive Behavioral Psychotherapies. *Cognitive Behavioral Therapy in Mental Health Care*, 249.
<https://www.torrossa.com/gs/resourceProxy?an=4913388&publisher=FZ7200#page=258>

Disciplined Personal Involvement. (2006). In J. P. McCullough, *Treating Chronic Depression with Disciplined Personal Involvement* (pp. 1–8). Springer US.
https://doi.org/10.1007/978-0-387-31066-4_1

Driscoll, K. A., Cukrowicz, K. C., Reardon, M. L., & Joiner, T. E. Jr. (2004). *Simple treatments for complex problems: A flexible cognitive behavior analysis system approach to psychotherapy* (2004-13620-000). Lawrence Erlbaum Associates Publishers.

Dunner, D. L. (2000). New Strategies for Improving the Treatment of Depression. *J Clin Psychiatry*, 61, 11. <https://www.psychiatrist.com/read-pdf/29670/>

Dunner, D. L. (2001). Acute and maintenance treatment of chronic depression. *Journal of Clinical Psychiatry*, 62, 10–16. https://www.psychiatrist.com/wp-content/uploads/2021/02/10425_acute-maintenance-treatment-chronic-depression.pdf

Elsaesser, M., Furukawa, T. A., & Schramm, E. (2022). Why one answer is not always enough: Reply to “CBASP may not be superior to other treatments for chronic depression”. *Journal of Affective Disorders*, 312, 144–145. <https://doi.org/10.1016/j.jad.2022.06.033>

Favorite, T. K. (n.d.). *The Use of CBASP in the Treatment of Chronic Depression and Co-morbid PTSD*. Retrieved April 24, 2024, from https://www.researchgate.net/profile/Todd-Favorite/publication/300961363_Behandlung_chronischer_Depression_und_komorbider_postraumatischer_Belastungsstorung_mit_dem_CBASP/links/5d442c7ba6fdcc370a74bc57/Behandlung-chronischer-Depression-und-komorbider-posttraumatischer-Belastungsstoerung-mit-dem-CBASP.pdf

Fosha, D. (2004). Brief Integrative Therapy Comes of Age: A Commentary. *Journal of Psychotherapy Integration*, 14(1), 66–92. <https://doi.org/10.1037/1053-0479.14.1.66>

Friedman, E. S., & Thase, M. E. (2007). DEPRESSION-FOCUSED. *Gabbard's Treatments of Psychiatric Disorders*, 409.
https://books.google.com/books?hl=en&lr=lang_en|lang_it&id=-T2aEqfwLW4C&oi=fnd&pg=PA409&dq=CBASP+or+cognitive+behavioral+analysis+system+of+psychotherapy+&ots=PCBoso-A9F&sig=mMvkaz707DIGms359fXlbfnxow

Gilbert Pinard, M. D. (n.d.). *New Developments in Cognitive-Behavioral Therapy for Depression*. Retrieved April 24, 2024, from <https://www.academia.edu/download/45385295/pinard-SF.pdf>

Guhn, A., & Brakemeier, E.-L. (2023). Psychotherapie: CBASP. In *Praxishandbuch Depression* (pp. 237–247). Elsevier.

Habtewold, T. D., van Bronswijk, S., Peeters, F., & Baştürk, N. (2022). Cognitive behavioral analysis system of psychotherapy (CBASP) may not be superior to other treatments for chronic depression: Correspondence. *Journal of Affective Disorders*, 308, 188–189. <https://doi.org/10.1016/j.jad.2022.04.049>

Hellerstein, D. J. (2008). *Review of Treating chronic depression with disciplined personal involvement: Cognitive behavioral analysis system of psychotherapy (CBASP)*. <https://psycnet.apa.org/record/2008-13167-013>

Hull, A., Swan, J., & McVicar, R. (2009). Treating Chronic Depression with Cognitive Behavioral Analysis System of Psychotherapy (CBASP). *European Psychiatry*, 24(S1), 1–1. <https://www.cambridge.org/core/journals/european-psychiatry/article/treating-chronic-depression-with-cognitive-behavioural-analysis-system-of-psychotherapy-cbasp/CFE6F8B027563C0155ABE04FA1E37912>

Kahl, K. G., Winter, L., & Schweiger, U. (2012). The third wave of cognitive behavioral therapies: What is new and what is effective? *Current Opinion in Psychiatry*, 25(6), 522–528. <https://doi.org/10.1097/YCO.0b013e328358e531>

Kahl, K. G., Winter, L., Schweiger, U., & Sipos, V. (2011). The third wave of cognitive-behavioral psychotherapies: Concepts and efficacy. *Fortschritte Der Neurologie-Psychiatrie*, 79(6), 330–339. <https://doi.org/10.1055/s-0029-1245963>

Kasnetz, M. D., McCullough, J. P., & Kaye, A. L. (1995). Patient Manual for Cognitive Behavioral Analysis System of Psychotherapy (CBASP). *Richmond: Virginia Commonwealth University*.

Keitner, G. I., & Cardemil, E. V. (2003). Psychotherapy for chronic depressive disorders. In *Handbook of Chronic Depression* (pp. 181–204). CRC Press.

<https://www.taylorfrancis.com/chapters/edit/10.4324/9780203912775-11/psychotherapy-chronic-depressive-disorders-gabor-keitner-esteban-cardemil>

Kocsis, J. H. (2000). New strategies for treating chronic depression. *Journal of Clinical Psychiatry*, 61, 42–45. <https://www.psychiatrist.com/read-pdf/20055/>

Köhler, S., Guhn, A., Betzler, F., Stiglmayr, C., Brakemeier, E.-L., & Sterzer, P. (2017). Therapeutic self-disclosure within DBT, schema therapy, and CBASP: Opportunities and challenges. *Frontiers in Psychology*, 8. <https://doi.org/10.3389/fpsyg.2017.02073>

Kramer, U., Belz, M., & Caspar, F. (2013). Psychotherapy of chronic depression: Contributions of CBASP by McCullough. *L'Encephale*, 39(2), 137–142. <https://doi.org/10.1016/j.encep.2012.03.006>

Lande, R. G. (2004). Review of Treatment for Chronic Depression: Cognitive Behavioral Analysis System of Psychotherapy. *Journal of Psychiatry & Law*, 32(2), 235–238. <http://proxy.library.stonybrook.edu/login?url=https://search.ebscohost.com/login.aspx?direct=true&db=psyh&AN=2004-20256-008&site=ehost-live&scope=site>

Long, L. R. (2023). *Dismantling Cognitive Behavioral Analysis System of Psychotherapy: A Pilot Study to Identify the Active Ingredients*. <https://digitalcommons.liberty.edu/doctoral/4263/>

Managing chronic depression: Harvard Mental Health Letter. (2009). *Harvard Mental Health Letter*, 26(6), 1–3. <http://proxy.library.stonybrook.edu/login?url=https://search.ebscohost.com/login.aspx?direct=true&db=pbh&AN=45142208&site=ehost-live&scope=site>

Matthews, J. D., & Doorley, J. (2016). Cognitive behavioral therapies for chronic depression. In T. J. Petersen, S. E. Sprich, & S. Wilhelm (Eds.), *The Massachusetts General Hospital handbook of cognitive behavioral therapy*. (2015-50703-022; pp. 301–317). Humana Press/Springer Nature. https://doi.org/10.1007/978-1-4939-2605-3_22

McCullough Jr, J. P. (2008). The cognitive behavioral analysis system of psychotherapy: A value-added strategy for chronic depression. *Psychiatric Times*, 25(10), 26–26.
<https://go.gale.com/ps/i.do?id=GALE%7CA184379074&sid=googleScholar&v=2.1&it=r&linkaccess=abs&issn=08932905&p=AONE&sw=w>

McCullough Jr, J. P. (2019). *Swimming Upstream: A story about becoming human*. Dorrance Publishing.
https://books.google.com/books?hl=en&lr=lang_en|lang_it&id=KKDHDwAAQBAJ&oi=fnd&pg=PT7&dq=CBASP+or+cognitive+behavioral+analysis+system+of+psychotherapy+&ots=uEvvSeJ-xm&sig=O0e6RSHhA1voAyvfpTiLtECBR1g

McCullough Jr., J. P. (2003). Treatment for chronic depression using Cognitive Behavioral Analysis System of Psychotherapy (CBASP). *Journal of Clinical Psychology*, 59(8), 833–846. <https://doi.org/10.1002/jclp.10176>

McCullough, J. P. (1995). Manual for Cognitive Behavioral Analytic System of Psychotherapy (CBASP). Richmond: Virginia Commonwealth University.

McCullough, J. P. Jr. (2000). *Treatment for chronic depression: Cognitive Behavioral Analysis System of Psychotherapy (CBASP)* (1999-04419-000). Guilford Press.

McCullough, J. P. Jr. (2001). *Skills training manual for diagnosing and treating chronic depression: Cognitive behavioral analysis system of psychotherapy* (2001-05096-000). Guilford Press.

McCullough, J. P. Jr. (2003). Treatment for chronic depression: Cognitive behavioral analysis system of psychotherapy (CBASP). *Journal of Psychotherapy Integration*, 13(3–4), 241–263. <https://doi.org/10.1037/1053-0479.13.3-4.241>

McCullough, J. P. Jr. (2005). Cognitive Behavioral Analysis System of Psychotherapy (CBASP) for Chronic Depression. In J. C. Norcross & M. R. Goldfried (Eds.), *Handbook of psychotherapy integration.*, 2nd ed. (2005-03070-013; pp. 281–298). Oxford University Press. <https://doi.org/10.1093/med:psych/9780195165791.003.0013>

McCullough, J. P. Jr. (2006a). Chronic Depression and the Cognitive Behavioral Analysis System of Psychotherapy. In G. Stricker & J. Gold (Eds.), *A casebook of psychotherapy integration*. (2006-03275-011; pp. 137–151). American Psychological Association.
<https://doi.org/10.1037/11436-011>

McCullough, J. P. Jr. (2006b). *Treating chronic depression with disciplined personal involvement* (2006-11486-000). Springer Science + Business Media.

McCullough, J. P. Jr. (2021). Characteristics of the optimal cognitive behavioral analysis system of psychotherapy (CBASP) therapist role. *Frontiers in Psychiatry*, 11.
<https://doi.org/10.3389/fpsy.2020.609954>

McCullough, J. P. Jr., & Schramm, E. (2019). Cognitive behavioral analysis system of psychotherapy for chronic depression. In J. C. Norcross & M. R. Goldfried (Eds.), *Handbook of psychotherapy integration.*, 3rd ed. (2019-28355-014; pp. 303–321). Oxford University Press. <https://doi.org/10.1093/med-psych/9780190690465.003.0014>

McCullough, J. P. Jr., Schramm, E., & Penberthy, J. K. (2015). *CBASP as a distinctive treatment for persistent depressive disorder* (2015-03274-000). Routledge/Taylor & Francis Group.

McGovern, C. (2024). Mark Berthold-Losleben, Marianne Liebing-Wilson, and John S. Swan: The ABCs of CBASP: A Guide to the Cognitive Behavioral Analysis System of Psychotherapy for Therapists and Supervisors: Hogrefe Publishing Corporation, Newburyport, MA, 2023, 196 pages, 11 figures, 4 tables, 10 boxes. *Journal of Contemporary Psychotherapy*, 54(1), 83–84. <https://doi.org/10.1007/s10879-023-09593-8>

McMahon, K., Herr, N. R., Zerubavel, N., Hoertel, N., & Neacsu, A. D. (2016). Psychotherapeutic treatment of bipolar depression. *Psychiatric Clinics*, 39(1), 35–56. [https://www.psych.theclinics.com/article/S0193-953X\(15\)00100-8/abstract](https://www.psych.theclinics.com/article/S0193-953X(15)00100-8/abstract)

Melrose, S. (2019). Persistent depressive disorder or dysthymia: An overview of assessment and treatment approaches. *Sherri Melrose Publications: A Virtual Memory Box*. <https://pressbooks.pub/sherrimelrosepublications/chapter/persistent-depressive-disorder-or-dysthymia-an-overview-of-assessment-and-treatment-approaches/>

Neudeck, P., Schoepf, D., & Penberthy, J. K. (2010). Learning theory aspects of the interpersonal discrimination exercise in cognitive behavioral analysis system of psychotherapy. *The Behavior Therapist*, 33(3), 58–63.

Neudeck, P., Walter, H., & Schoepf, D. (2012). Exposure aspects of the interpersonal discrimination exercise (IDE) and the situational analysis (SA) in cognitive behavioral analysis system of psychotherapy (CBASP). In P. Neudeck & H.-U. Wittchen (Eds.), *Exposure therapy: Rethinking the model—Refining the method*. (2012-16171-009; pp. 153–166). Springer Science + Business Media. https://doi.org/10.1007/978-1-4614-3342-2_9

Newman, C. F. (2013). *Core competencies in cognitive-behavioral therapy: Becoming a highly effective and competent cognitive-behavioral therapist*. Routledge. <https://www.taylorfrancis.com/books/mono/10.4324/9780203107447/core-competencies-cognitive-behavioral-therapy-cory-newman>

Norcross, J. C., & Goldfried, M. R. (2005). *Handbook of psychotherapy integration*. Oxford University Press. https://books.google.com/books?hl=en&lr=lang_en|lang_it&id=w-wpACyzL9gC&oi=fnd&pg=PR9&dq=CBASP+or+Cognitive+Behavioral+Analysis+System+of+Psychotherapy+&ots=QKH0suglb_&sig=KvkqcgwuN1ElaaYMho9RTT7zkgM

Oertel-Knöchel, V., & Schoepf, D. (2013). Adaptation of cognitive behavioral analysis system of psychotherapy in a 29 year old female patient with Chronic Major Depression and antecedent dysthymic disorder who switched under combined SSRI/CBASP outpatient treatment into bipolarity: A case report. In *Psychiatric Disorders-New Frontiers in Affective Disorders*. IntechOpen. <https://www.intechopen.com/chapters/44830>

Penberthy, J. K. (2019). *Persistent Depressive Disorder* (Vol. 43). Hogrefe Publishing GmbH. https://books.google.com/books?hl=en&lr=lang_en|lang_it&id=0pbEAAAQBAJ&coi=fnd&pg=PR5&dq=CBASP+or+cognitive+behavioral+analysis+system+of+psychotherapy+&ots=v-et CSmjR&sig=SKB1pJ_WCnPj4Zpj5ocDuJlbJFM

Pettit, J. W., & Joiner, T. E. (2006). *Therapeutics*. American Psychological Association. <https://psycnet.apa.org/record/2005-11928-013>

Ramsay, J. R. (2001). Review of Treatment for chronic depression: Cognitive behavioral analysis system of psychotherapy. *Journal of Psychotherapy Integration*, 11(3), 415–419. <https://doi.org/10.1023/A:1016662402537>

Riso, L. P. (2002). Introduction: Considerations for Psychotherapy with Chronically Depressed Patients. *Journal of Contemporary Psychotherapy*, 32(4), 259. <https://search.proquest.com/openview/b05a8b50a4bfae8a6bc9b1dbfc57def/1?pq-origsite=gscholar&cbl=48134>

Riso, L. P., & Thase, M. E. (2007). Chronic Depression. In N. Kazantzis & L. L'Abate (Eds.), *Handbook of homework assignments in psychotherapy: Research, practice, prevention*. (2006-11928-014; pp. 247–262). Springer Science + Business Media. https://doi.org/10.1007/978-0-387-29681-4_15

Riso, L. P., McCullough, J. P., & Blandino, J. (2003). Focus on Empirically Supported Methods: The Cognitive Behavioral Analysis System of Psychotherapy: A Promising Treatment for Clinical Depression. *The Scientific Review of Mental Health Practice: Objective Investigations of Controversial and Unorthodox Claims in Clinical Psychology, Psychiatry, and Social Work*, 2(1), 61–68.

Rose, F. E. (2000). Treatment for chronic depression: Cognitive behavioral analysis system of psychotherapy (CBASP). *Depression and Anxiety*, 12(1), 56–57. [https://doi.org/10.1002/1520-6394\(2000\)12:1<56::AID-DA10>3.0.CO;2-U](https://doi.org/10.1002/1520-6394(2000)12:1<56::AID-DA10>3.0.CO;2-U)

Ruggiero, G. M., Spada, M. M., Caselli, G., & Sassaroli, S. (2018). A Historical and Theoretical Review of Cognitive Behavioral Therapies. *Journal of Rational-Emotive & Cognitive-Behavior Therapy*, 36(4), 378–403. <https://doi.org/10.1007/s10942-018-0292-8>

Salomon, R. M. (2000). *Treatment for Chronic Depression: Cognitive Behavioral Analysis System of Psychotherapy (CBASP)*. Physicians Postgraduate Press, Inc. <https://www.psychiatrist.com/jcp/treatment-chronic-depression-cognitive-behavioral/>

Salton, J. A. (2007). *Treating chronic depression with disciplined personal involvement: Cognitive Behavioral Analysis System of Psychotherapy (CBASP)*. SciELO Brasil.
<https://www.scielo.br/j/rprs/a/hnG8j6qstc9MmP7QNwCdjdv/?lang=en>

Sayegh, L. (n.d.). *GROUP TREATMENT MANUAL FOR PERSISTENT DEPRESSION*. Retrieved April 24, 2024, from
<http://ndl.ethernet.edu.et/bitstream/123456789/55282/1/35.pdf>

Sayegh, L., & Penberthy, J. K. (2016). *Group Workbook for Treatment of Persistent Depression: Cognitive Behavioral Analysis System of Psychotherapy-(CBASP) Patient's Guide*. Routledge.
<https://www.taylorfrancis.com/books/mono/10.4324/9781315674544/group-workbook-treatment-persistent-depression-liliane-sayegh-kim-penberthy>

Sayegh, L., & Turecki, G. (2013). Group Cognitive Behavioral Analysis System of Psychotherapy (Group-CBASP). *Psychiatric Disorders—New Frontiers in Affective Disorders*, 203.
https://books.google.com/books?hl=en&lr=lang_en|lang_it&id=NkueDwAAQBAJ&oi=fnd&pg=PA203&dq=CBASP+or+Cognitive+Behavioral+Analysis+System+of+Psychotherapy+&ots=0NBdahPKOH&sig=0NBqB2ZcY2xogfCNKpzYrQ4crow

Schramm, E., Caspar, F., & Berger, M. (2006). A specific therapy for chronic depression. McCullough's Cognitive Behavioral Analysis System of Psychotherapy. *Der Nervenarzt*, 77(3), 355–370; quiz 371. <https://doi.org/10.1007/s00115-006-2059-1>

Serbanescu, I.-D. (2023). *Towards Personalized Medicine for Persistent Depressive Disorder: Moving from “one size fits all” to “what works for whom?”* [PhD Thesis].
<https://archiv.ub.uni-heidelberg.de/volltextserver/33283/>

Sisemore, T. A. (2005). Review of Simple Treatments for Complex Problems: A Flexible Cognitive Behavioral Analysis System Approach To Psychotherapy. *Journal of Psychology and Christianity*, 24(2), 177–178.

Sperry, L. (2009). Selecting Psychotherapeutic Interventions. *Journal of Individual Psychology*, 65(3).
<https://search.ebscohost.com/login.aspx?direct=true&profile=ehost&scope=site&authtype=crawler&jrnl=15222527&AN=46739076&h=MutUfsbyRdeNnTF8WuwGakA9Wz105OT6CsbnfRmTmpOV66OvEhodNHanBkZ5P7GTHUjOIMrWRjtmiYbKdo1eAw%3D%3D&crl=c>

Sperry, L. (2016). *Handbook of diagnosis and treatment of DSM-5 personality disorders: Assessment, case conceptualization, and treatment*. Routledge.
<https://www.taylorfrancis.com/books/mono/10.4324/9780203763728/handbook-diagnosis-treatment-dsm-5-personality-disorders-len-sperry-len-sperry>

Swan, J. S., & Hull, A. M. (2007). The cognitive behavioural analysis system of psychotherapy: A new psychotherapy for chronic depression. *Advances in Psychiatric Treatment*, 13(6), 458–469. <https://www.cambridge.org/core/journals/advances-in-psychiatric-treatment/article/cognitive-behavioural-analysis-system-of-psychotherapy-a-new-psychotherapy-for-chronic-depression/DDD5575ADC4425D7BC548D4D3104005B>

Swan, J., Liebing-Wilson, M., & Sloan, G. (2015a). An introduction to the cognitive behavioral analysis system of psychotherapy: Part 2. *British Journal of Mental Health Nursing*, 4(6), 260–266. <https://doi.org/10.12968/bjmh.2015.4.6.260>

Swan, J., Liebing-Wilson, M., & Sloan, G. (2015b). The cognitive behavioral analysis system of psychotherapy. *British Journal of Mental Health Nursing*, 4(5), 206–212. <https://doi.org/10.12968/bjmh.2015.4.5.206>

Treating the Chronically Depressed Patient. (2006). In J. P. McCullough, *Treating Chronic Depression with Disciplined Personal Involvement* (pp. 38–53). Springer US. https://doi.org/10.1007/978-0-387-31066-4_3

Uher, R. (2014). Persistent depressive disorder, dysthymia, and chronic depression: Update on diagnosis, treatment. *Psychiatric Times*, 31(8), 46–46. <https://go.gale.com/ps/i.do?id=GALE%7CA377664540&sid=googleScholar&v=2.1&it=r&linkaccess=abs&issn=08932905&p=AONE&sw=w>

Vivian, D. (2019). Principles of Therapeutic Change in Treating Depression with an Integrative Application of the Cognitive Behavioral Analysis System of Psychotherapy. *Principles of Change: How Psychotherapists Implement Research in Practice*, 129. https://books.google.com/books?hl=en&lr=lang_en|lang_it&id=SCeeDwAAQBAJ&oi=fnd&pg=PA129&dq=CBASP+or+Cognitive+Behavioral+Analysis+System+of+Psychotherapy+&ots=Mi7hBO3X_9&sig=W3-Q19XuQf1YsdrMbTnlOfLUeHw

Vivian, D., & Salwen, J. (2013). Key process issues in Cognitive Behavioral Analysis System of Psychotherapy (CBASP): Translation of an evidence-based model into clinical practice and training. *Psychotherapy*, 50(3), 398–403. <https://doi.org/10.1037/a0033054>

Wiersma, J. E., van Schaik, D. J. F., Blom, M. B. J., Bakker, L., van Oppen, P., & Beekman, A. T. F. (2009). Treatment for chronic depression: Cognitive behavioral analysis system of psychotherapy (CBASP). *Tijdschrift Voor Psychiatrie*, 51(10), 727–736.

Wiersma, J., Klein, P., Schramm, E., Furukawa, T., & Favorite, T. (2021). Editorial: CBASP in the Treatment of Persistent Depressive Disorder. *Frontiers in Psychiatry*, 12, 804602. <https://doi.org/10.3389/fpsyg.2021.804602>

Williams, M. J., & Jarrett, R. B. (2005). Depression-Adult. In A. Freeman, S. H. Felgoise, C. M. Nezu, A. M. Nezu, & M. A. Reinecke (Eds.), *Encyclopedia of Cognitive Behavior Therapy* (pp. 151–155). Springer-Verlag. https://doi.org/10.1007/0-306-48581-8_45

Non-English Literature
2008-2022 (8)

Brakemeier, E.-L. (2022). Psychotherapie – Ein kompetenzorientiertes Lehrbuch. *Psychotherapeut*, 67(1), 81–84. <https://doi.org/10.1007/s00278-021-00556-9>

Gruppenpsychotherapie, I., & kognitive Therapie, A. (2012). Depressive Störungen. *Gruppenpsychotherapie: Lehrbuch Für Die Praxis*, 259.
https://books.google.com/books?hl=en&lr=lang_en|lang_it&id=yOE3DwAAQBAJ&oi=fnd&pg=PA255&dq=CBASP+or+cognitive+behavioral+analysis+system+of+psychotherapy+&ots=b7Ndl8SJ4D&sig=t9syMhgeIyG2_d3psd2CR1Lpzpw

Hohagen, F., & Lotz-Rambaldi, W. (2011). Zur fragmentierung in der verhaltenstherapie. *Verhaltenstherapie*, 21(2), 84–85.
<https://scholar.archive.org/work/gxvftzbaxfgubmr75ml2it3dma/access/wayback/https://www.karger.com/Article/Pdf/329404>

Hohagen, F., & Lotz-Rambaldi, W. (2012). Plädoyer für eine modulare Psychotherapie. *Verhaltenstherapie*, 22(2), 84–85.
<https://scholar.archive.org/work/ajj5sihhavfgthz4nh5qefu7fm/access/wayback/https://www.karger.com/Article/Pdf/339443>

Kahl, K. G., Winter, L., Schweiger, U., & Sipos, V. (2011). The third wave of cognitive-behavioural psychotherapies: Concepts and efficacy. *Fortschritte Der Neurologie-Psychiatrie*, 79(6), 330–339. <https://doi.org/10.1055/s-0029-1245963>

Klein, J. P., Steinlechner, S., Sipos, V., & Schweiger, U. (2011). Psychotherapy of chronic depression with cognitive behavioral analysis system of psychotherapy (CBASP). *Psychotherapie, Psychosomatik, Medizinische Psychologie*, 61(12), 526–534; quiz 535.
<https://doi.org/10.1055/s-0031-1276990>

Teismann, T., Michalak, J., Willutzki, U., Walter, D., Boyraz, M., Döpfner, M., Linden, M., Tominschek, I., Schiepek, G., & Mehl, C. (2008). James McCullough: «Die Rolle des Therapeuten im CBASP ist die eines aktiven Mitspielers». *Verhaltenstherapie*, 18(3), 186–188. <https://karger.com/ver/article-abstract/18/3/186/319183>

Wambach, K., & Rief, W. (2012). Ist Cognitive Behavioral Analysis System of Psychotherapy (CBASP) eine Verhaltenstherapie? *Verhaltenstherapie*, 22(4), 224–226.
https://www.researchgate.net/profile/Katrin-Wambach/publication/270724659_Is_Cognitive_Behavioral_Analysis_System_of_Psychotherapy_CBASP_Behavioral_Therapy/links/56002ae08aeafc8ac8c4e60/Is-Cognitive-Behavioral-Analysis-System-of-Psychotherapy-CBASP-Behavioral-Therapy.pdf